Pounce™ Thrombectomy System | Endologix

CASE REVIEW

Brachial Embolic Occlusion

Case Demographics

Age (years): 84
Gender: Female

Clot Removal

Pounce™ Thrombectomy System
(no thrombolysis or aspiration)

Clinical Presentation


- Sudden onset of cold, painful right upper extremity
- Hx of chronic atrial fibrillation (anticoagulation had been temporarily discontinued)
- Angiography revealed a tortuous brachial artery with embolus obstructing the radial and ulnar arteries

Procedural Information


Intervention took place promptly after Dx. One pass was made with the Pounce Thrombectomy System, removing the embolus in the brachial artery and reestablishing flow into radial and ulnar arteries. No thrombolysis was used.

Anatomy

Pre



Post



INTENDED USE: The Pounce™ Thrombectomy System is intended for the non-surgical removal of thrombi and emboli from the peripheral arterial vasculature.

CONTRAINDICATIONS:

- The device is contraindicated for use in patients who cannot receive recommended intravenous anticoagulant therapy.
- The device is not for coronary or neurovascular use.
- The device is not intended to be deployed in vessels with previously implanted devices.

Refer to Instructions for Use for more information concerning Indications, Contraindications, Specific Anatomic Considerations, Warnings, Precautions, and Adverse Events.

CAUTION: Federal (USA) law restricts this device to sale by or on the order of a physician.

NOTE: Not all product components are available in every country.

Please consult with your Endologix representative to confirm product availability.

Endologix, Pounce and the Endologix and Pounce logos are trademarks of Endologix LLC and/or its affiliates. All other trademarks are the respective property of their owners.
© 2026 Endologix LLC. All rights reserved. MM3064-US Rev 01